

ASTRAZENECA PLC  
Form 6-K  
January 12, 2018

FORM 6-K

SECURITIES AND EXCHANGE COMMISSION  
Washington, D.C. 20549

Report of Foreign Issuer

Pursuant to Rule 13a-16 or 15d-16 of  
the Securities Exchange Act of 1934

For the month of January 2018

Commission File Number: 001-11960

AstraZeneca PLC

1 Francis Crick Avenue  
Cambridge Biomedical Campus  
Cambridge CB2 0AA  
United Kingdom

Indicate by check mark whether the registrant files or will file annual reports under cover of Form 20-F or Form 40-F.

Form 20-F ☒ Form 40-F ☐

Indicate by check mark if the registrant is submitting the Form 6-K in paper as permitted by Regulation S-T Rule 101(b)(1):

Indicate by check mark if the registrant is submitting the Form 6-K in paper as permitted by Regulation S-T Rule 101(b)(7): ☐

Indicate by check mark whether the registrant by furnishing the information contained in this Form is also thereby furnishing the information to the Commission pursuant to Rule 12g3-2(b) under the Securities Exchange Act of 1934.

Yes ☐ No ☒

If "Yes" is marked, indicate below the file number assigned to the Registrant in connection with Rule 12g3-2(b):  
82- \_\_\_\_\_

12 January 2018 16:10 GMT

LYNPARZA APPROVED BY US FDA IN GERMLINE BRCA-MUTATED  
METASTATIC BREAST CANCER

Lynparza is the first and only PARP inhibitor approved for use beyond ovarian cancer

Lynparza reduced the risk of disease progression or death by 42% compared to standard of care chemotherapy

AstraZeneca and Merck & Co., Inc., Kenilworth, N.J., US (Merck: known as MSD outside the US and Canada) today announced that the US Food and Drug Administration (FDA) has approved Lynparza (olaparib), for use in patients with deleterious or suspected deleterious germline BRCA-mutated (gBRCAm), human epidermal growth factor receptor 2 (HER2)-negative metastatic breast cancer who have been previously treated with chemotherapy in the neoadjuvant, adjuvant or metastatic setting. Patients with hormone receptor positive (HR+) breast cancer should have been treated with a prior endocrine therapy or be considered inappropriate for endocrine therapy. Patients are selected for therapy based on an FDA-approved companion diagnostic from Myriad Genetics.

Dave Fredrickson, Executive Vice President, Head of the Oncology Business Unit, AstraZeneca, said: "This new approval for Lynparza makes it the first and only PARP inhibitor approved in metastatic breast cancer, and the only PARP inhibitor approved beyond ovarian cancer. This is significant for breast cancer patients, as the identification of BRCA status, in addition to hormone receptor and HER2 status, becomes a potentially critical step in the management of their disease."

Roy Baynes, Senior Vice President and Head of Global Clinical Development, Chief Medical Officer, Merck Research Laboratories said: "This additional approval for Lynparza represents an important advance for women with HER2-negative metastatic breast cancer with a germline BRCA mutation, which is a difficult-to-treat cancer. Moreover, this approval adds further impetus to our important collaboration with AstraZeneca in developing cancer therapies."

The approval was based on data from the randomised, open-label, Phase III OlympiAD trial which investigated Lynparza versus physician's choice of chemotherapy (capecitabine, eribulin, or vinorelbine). In the trial, Lynparza significantly prolonged progression-free survival (PFS) compared with chemotherapy, and reduced the risk of disease progression or death by 42% (HR 0.58; 95% CI 0.43-0.80; P=0.0009 median 7.0 vs 4.2 months). Patients with measurable disease taking Lynparza (n=167) experienced an objective response rate of 52% (95% CI 44-60), double the response rate for those in the chemotherapy arm (n=66) which was 23% (95% CI 13-35). Additionally, patients experienced a confirmed complete response rate of 7.8% for Lynparza compared to 1.5% for the chemotherapy arm. The data from the OlympiAD trial can be found in the June 2017 issue of the New England Journal of Medicine.

Susan M. Domchek, Executive Director of the Bassett Center for BRCA at the Abramson Cancer Center of the University of Pennsylvania, and a national leader on the OlympiAD trials, said: "Patients diagnosed with BRCA-related metastatic breast cancer are often younger than other breast cancer patients, and their disease is often much more aggressive and difficult to treat. While there is currently no cure for metastatic breast cancer, today's approval offers a new, targeted option that may help to delay disease progression for these patients."

The most common adverse reactions ( $\geq 20\%$ ) in the OlympiAD trial of patients who received Lynparza were nausea (58%), anaemia (40%), fatigue (including asthenia) (37%), vomiting (30%), neutropenia (27%), respiratory tract infection (27%), leukopenia (25%), diarrhoea (21%), and headache (20%). The percentage of patients who discontinued treatment in the Lynparza arm was 5% compared to the chemotherapy arm which was 8%.

This is the third indication approved for Lynparza in the US, where it has been used to treat nearly 4,000 advanced ovarian cancer patients. Lynparza has the broadest clinical development programme of any PARP inhibitor, and AstraZeneca and MSD are working together to deliver Lynparza as quickly as possible to more patients across multiple settings, including breast, ovarian, prostate and pancreatic cancers.

Sustainable and Ongoing Externalisation Revenue

Under the oncology collaboration with Merck, announced in July 2017, AstraZeneca is potentially eligible for more than \$6 billion of future Sustainable and Ongoing Externalisation Revenue in the form of sales-related and approval-related payments in addition to option payments until 2019. Following this new approval for Lynparza, AstraZeneca will receive \$70 million in Sustainable and Ongoing Externalisation Revenue.

#### About OlympiAD

OlympiAD is a randomised, open-label, multicentre Phase III trial assessing the efficacy and safety of Lynparza tablets (300 mg twice daily) compared to physician's choice of chemotherapy in 302 patients with HER2-negative metastatic breast cancer with germline BRCA1 or BRCA2 mutations, which are confirmed or suspected to be deleterious. The international trial was conducted in 19 countries across Europe, Asia, North America and South America.

Patients in the OlympiAD trial had HER2-negative gBRCA1- or gBRCA2-mutated breast cancer, which was HR+ or triple negative, and received Lynparza for metastatic disease. Approximately half of the patients in the Lynparza and chemotherapy arm of the trial were HR+ (n=152), and approximately half were triple negative (n=150). Among the 205 patients treated with Lynparza, the median age was 44 years (range: 22 to 76). Before enrolment, patients had prior treatment with an anthracycline (unless contraindicated) and a taxane chemotherapy either in the neoadjuvant, adjuvant or metastatic setting and no more than two prior lines of chemotherapy for metastatic disease. Hormone receptor-positive patients had received at least one endocrine medicine or were not eligible for endocrine medicines. Prior treatments with endocrine medicines were not counted as prior lines of chemotherapy.

The primary endpoint of the trial was PFS as measured by a Blinded Independent Central Review. Secondary endpoints included overall survival, time to second progression or death, objective response rate, and effect on health-related quality of life.

#### About Metastatic Breast Cancer (MBC)

Three main receptors drive tumour growth in breast cancer: progesterone receptors (PR), estrogen receptors (ER) and HER2 receptors. A patient's breast cancer will test either negative or positive for these three receptors. If a tumour tests positive for PR and/or ER, it is considered HR+. If a tumour tests negative for all three receptors, it is considered triple negative.

MBC is the most advanced stage of breast cancer (Stage IV), and occurs when cancer cells have spread beyond the initial tumour site to other parts of the body outside of the breast.

Despite the increase in treatment options during the past three decades, there is currently no cure for patients diagnosed with MBC and only 26.9% of patients survive five years after diagnosis. Thus, the primary aim of treatment is to slow progression of the disease for as long as possible, improving, or at least maintaining, a patient's quality of life.

It is estimated that in 2018, there will be approximately 155,000 women in the US living with MBC, and this number is projected to increase to approximately 160,000 by the year 2020.

#### About Germline BRCA Mutations

BRCA1 and BRCA2 are human genes that produce proteins responsible for repairing damaged DNA and play an important role maintaining the genetic stability of cells. When either of these genes is mutated, or altered, such that its protein product either is not made or does not function correctly, DNA damage may not be repaired properly and cells become unstable. As a result, cells are more likely to develop additional genetic alterations that can lead to cancer.

#### About Lynparza (olaparib)

Lynparza is the first FDA-approved oral poly ADP-ribose polymerase (PARP) inhibitor and the first targeted treatment to potentially exploit DNA damage response (DDR) pathway deficiencies, such as BRCA mutations, to

preferentially kill cancer cells. Specifically, in vitro studies have shown that Lynparza-induced cytotoxicity may involve inhibition of PARP enzymatic activity and increased formation of PARP-DNA complexes, resulting in DNA damage and cancer cell death.

Lynparza is being investigated in a range of DDR-deficient tumour types and is the foundation of AstraZeneca's industry-leading portfolio of compounds targeting DDR mechanisms in cancer cells.

#### About the AstraZeneca and MSD Strategic Oncology Collaboration

In July 2017, AstraZeneca and Merck & Co., Inc., Kenilworth, NJ, US, known as MSD outside the United States and Canada, announced a global strategic oncology collaboration to co-develop and co-commercialise Lynparza, the world's first PARP inhibitor, and potential new medicine selumetinib, a MEK inhibitor, for multiple cancer types. The collaboration is based on increasing evidence that PARP and MEK inhibitors can be combined with PD-L1/PD-1 inhibitors for a range of tumour types. Working together, the companies will develop Lynparza and selumetinib in combination with other potential new medicines and as a monotherapy. Independently, the companies will develop Lynparza and selumetinib in combination with their respective PD-L1 and PD-1 medicines.

#### About AstraZeneca in Oncology

AstraZeneca has a deep-rooted heritage in Oncology and offers a quickly growing portfolio of new medicines that has the potential to transform patients' lives and the Company's future. With at least six new medicines to be launched between 2014 and 2020 and a broad pipeline of small molecules and biologics in development, we are committed to advance New Oncology as one of AstraZeneca's five Growth Platforms focused on lung, ovarian, breast and blood cancers. In addition to our core capabilities, we actively pursue innovative partnerships and investments that accelerate the delivery of our strategy as illustrated by our investment in Acerta Pharma in haematology.

By harnessing the power of four scientific platforms - Immuno-Oncology, Tumour Drivers and Resistance, DNA Damage Response and Antibody Drug Conjugates - and by championing the development of personalized combinations, AstraZeneca has the vision to redefine cancer treatment and one day eliminate cancer as a cause of death.

#### About AstraZeneca

AstraZeneca is a global, science-led biopharmaceutical company that focuses on the discovery, development and commercialisation of prescription medicines, primarily for the treatment of diseases in three therapy areas - Oncology, Cardiovascular & Metabolic Diseases and Respiratory. The Company also is selectively active in the areas of autoimmunity, neuroscience and infection. AstraZeneca operates in over 100 countries and its innovative medicines are used by millions of patients worldwide.

For more information, please visit [www.astrazeneca.com](http://www.astrazeneca.com) and follow us on Twitter @AstraZeneca.

#### Media Relations

Esra Erkal-Paler	UK/Global	+44 203 749 5638
Karen Birmingham	UK/Global	+44 203 749 5634
Rob Skelding	UK/Global	+44 203 749 5821
Matt Kent	UK/Global	+44 203 749 5906
Gonzalo Viña	UK/Global	+44 203 749 5916
Jacob Lund	Sweden	

Edgar Filing: ASTRAZENECA PLC - Form 6-K

Michele Meixell	US	+46 8 553 260 20 +1 302 885 2677
Investor Relations		
Thomas Kudsk Larsen		+44 203 749 5712
Craig Marks	Finance, Fixed Income, M&A	+44 7881 615 764
Henry Wheeler	Oncology	+44 203 749 5797
Mitchell Chan	Oncology; Other	+1 240 477 3771
Christer Gruvris	Brilinta; Diabetes	+44 203 749 5711
Nick Stone	Respiratory; Renal	+44 203 749 5716
US toll free		+1 866 381 7277

Adrian Kemp  
Company Secretary  
AstraZeneca PLC

SIGNATURES

Pursuant to the requirements of the Securities Exchange Act of 1934, the Registrant has duly caused this report to be signed on its behalf by the undersigned, thereunto duly authorized.

AstraZeneca PLC

Date: 12 January 2018

By: /s/ Adrian Kemp  
Name: Adrian Kemp  
Title: Company Secretary